



“Baseline characteristics of idiopathic pulmonary fibrosis: analysis from the Australian Idiopathic Pulmonary Fibrosis Registry.” Helen E. Jo, Ian Glaspole, Christopher Grainge, Nicole Goh, Peter M.A. Hopkins, Yuben Moodley, Paul N. Reynolds, Sally Chapman, E. Haydn Walters, Christopher Zappala, Heather Allan, Gregory J. Keir, Andrew Hayen, Wendy A. Cooper, Annabelle M. Mahar, Samantha Ellis, Sacha Macansh and Tamera J. Corte. *Eur Respir J* 2017; 49: 1601592.

This article from the February 2017 issue of the *European Respiratory Journal* was published with an error in figure 5. The legend identifying “Anti-fibrotics” and “No anti-fibrotics” in figure 5c was mistakenly transposed in production. The figure has been corrected and is reproduced below. The article has been corrected and republished online.




FIGURE 5 a) Kaplan-Meier analysis for mortality for the Australian Idiopathic Pulmonary Fibrosis Registry cohort. b) Kaplan-Meier analysis according to GAP stage. GAP: gender, age, physiology. Log rank test  $p<0.0001$ . c) Kaplan-Meier analysis by use of anti-fibrotic treatment. Log rank test  $p<0.0001$ .

Copyright ©ERS 2017