



Perspectives of patients and healthcare professionals on mHealth for asthma self-management

Andrew J. Simpson¹, Persijn J. Honkoop², Erika Kennington³,
Jiska B. Snoeck-Stroband², Ian Smith², Jessica East³, Courtney Coleman³,
Ann Caress¹, Kian Fan Chung⁴, Jacob K. Sont², Omar Usmani⁴ and
Stephen J. Fowler¹

Affiliations: ¹Division of Infection, Immunity and Respiratory Medicine, School of Biological Sciences, The University of Manchester and University Hospital of South Manchester NHS Foundation Trust, Manchester, UK. ²Dept of Medical Decision Making, Leiden University Medical Center, Leiden, the Netherlands. ³Asthma UK, London, UK. ⁴National Heart and Lung Institute, Imperial College London and NIHR Biomedical Research Unit, Royal Brompton and Harefield NHS Trust, London, UK.

Correspondence: Andrew J. Simpson, Division of Infection, Immunity and Respiratory Medicine, University Hospital of South Manchester, Education and Research Centre (2nd Floor), Southmoor Road, Manchester M23 9LT, UK. E-mail: Andrew.Simpson-2@manchester.ac.uk



@ERSpublications

People with asthma and healthcare professionals strongly support the utility of mHealth for asthma self-management <http://ow.ly/wyJC3096f4e>

Cite this article as: Simpson AJ, Honkoop PJ, Kennington E, *et al.* Perspectives of patients and healthcare professionals on mHealth for asthma self-management. *Eur Respir J* 2017; 49: 1601966 [https://doi.org/10.1183/13993003.01966-2016].

ABSTRACT Mobile healthcare (mHealth) has the potential to revolutionise the self-management of long-term medical conditions such as asthma. A user-centred design is integral if mHealth is to be embraced by patients and healthcare professionals.

The aim of this study was to determine the perspectives of individuals with asthma and healthcare professionals on the use of mHealth for asthma self-management.

We used a sequential exploratory mixed methods design; focus groups informed the development of questionnaires, which were disseminated to individuals with asthma and healthcare professionals.

Focus group participants (18 asthma patients and five healthcare professionals) identified 12 potential uses of mHealth. Questionnaire results showed that individuals with asthma (n=186) most frequently requested an mHealth system to monitor asthma over time (72%) and to collect data to present to healthcare teams (70%). In contrast, healthcare professionals (n=63) most frequently selected a system alerting patients to deteriorating asthma control (86%) and advising them when to seek medical attention (87%). Individuals with asthma were less likely than healthcare professionals ($p<0.001$) to believe that assessing medication adherence and inhaler technique could improve asthma control.

Our data provide strong support for mHealth for asthma self-management, but highlight fundamental differences between the perspectives of patients and healthcare professionals.

Received: Oct 06 2016 | Accepted after revision: Feb 07 2017

Support statement: This project has received funding from the European Union's Horizon 2020 Framework Programme for Research and Innovation under grant agreement No. 643607. The funder had no input in the study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. The University Hospital of South Manchester NHS Trust sponsored this study. The sponsor had no input in the study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. Funding information for this article has been deposited with the Crossref Funder Registry.

Conflict of interest: Disclosures can be found alongside this article at erj.ersjournals.com

Copyright ©ERS 2017

Introduction

Asthma affects approximately 300 million people worldwide [1]. In many cases asthma control remains suboptimal and avoidable deaths are still occurring [2]. Asthma self-management has beneficial health outcomes such as reduced hospital admissions, better lung function, fewer asthma symptoms and less use of rescue medication [3–6]. Asthma guidelines recommend that all people with asthma receive education on asthma self-management [7, 8].

Traditional asthma self-management programmes utilise personal asthma action plans, which involve the monitoring of symptoms and/or peak flow, with a written action plan detailing how to recognise and respond to worsening asthma. This “pen and paper” approach is burdensome and time-consuming and neither patients nor healthcare teams are enthusiastic about its use [9]. Furthermore, action points based on rudimentary data, such as symptoms and peak flow, may be less effective than action points based on multiple personalised parameters [10]. Despite important benefits of self-management, only 27% of adults with asthma receive an asthma action plan [11] and patients’ adherence to written action plans is poor [12].

Web-based systems offer less burdensome self-management support, which may improve asthma outcomes [13]. Smartphones have become an integral part of life and mobile healthcare (mHealth) systems are promising tools that could revolutionise asthma self-management. There are over 200 mobile phone applications for asthma [14] and supplementary wearable and inhaler-based devices are widely available [15]. Currently, however, the utility of mHealth for asthma self-management is unknown and a recent Cochrane review was unable to advise clinicians and the general public on their efficacy [16]. Investigations into successful mHealth systems point to user-centred design practices [17]. In terms of asthma self-management, however, there is little evidence of user-centred design practices and no data exploring the perspectives of both end-users (*i.e.* patients and healthcare professionals).

The current investigation is part of the myAirCoach project, supported by the European Union’s Horizon 2020 Framework Programme for Research and Innovation. This project aims to create a user-centred mHealth tool to support the self-management of asthma. Understanding patients’ and healthcare professionals’ perspectives is a fundamental step in the development of user-centred mHealth systems. Therefore, the present study was conducted to determine the perspectives of individuals with asthma and healthcare professionals on the use of mHealth systems to support asthma self-management. Specifically, we aimed to determine end-users’: 1) experiences and perceived uses of mHealth systems for asthma self-management; 2) views of what measurements would be useful in managing asthma; and 3) perspectives on the acceptability of and barriers to using mHealth systems for asthma self-management.

Methods

Study design

We employed a sequential exploratory mixed methods design [18], in which qualitative exploration (using focus groups) informed instrument development for a subsequent electronic questionnaire, with findings from both data sources integrated.

Focus groups

One moderator guided each focus group, following the approach from Greenbaum [19], according to a structured schedule of topics (table 1); the content of which was determined *via* consultation with the myAirCoach collaborators (www.myaircoach.eu) and a patient advisory group. Three focus groups were conducted with individuals with asthma, in Manchester (UK), London (UK) and Leiden (the Netherlands), and one focus group was conducted with healthcare professionals in Manchester (UK). The focus groups were video-recorded, transcribed, translated where appropriate and underwent Framework Analysis [20]. Data were grouped under emergent themes and integrated into three predetermined core categories relating to the research aims. Data management was supported by NVivo qualitative analysis software (Version 10).

Individuals aged ≥ 18 years who either had doctor-diagnosed asthma or were healthcare professionals involved in the treatment of asthma were eligible to participate in the focus groups. Patients were recruited from respiratory clinics in Manchester and Leiden, and *via* advertisements placed online at Asthma UK’s website. Healthcare professionals were recruited from hospital respiratory departments in Manchester.

Questionnaires

A long-list of questions was generated following analysis of the focus group transcripts. The clinical research team and members of the hospital’s patient and public involvement team (including a patient representative) assessed the questions for face validity and bias. The importance of each question was then ranked and the number of items reduced to generate a 20-item questionnaire for people with asthma and a 10-item questionnaire for healthcare professionals. The questionnaires were different for people with asthma and healthcare professionals, but contained some identical questions to allow comparisons between

TABLE 1 Focus group topic guide for people with asthma and healthcare professionals

Topic 1. Experiences and perceived uses of mHealth for asthma

Example prompts: Have you previously used mHealth systems to help manage your/your patients' asthma? What would you consider would be a useful purpose of an mHealth system with regards to your/your patients' asthma?

Topic 2. Potential useful measurements for mHealth

Example prompts: What physiological, behavioural and environmental measurements could help you manage your/your patients' asthma?

Topic 3. Burden and barriers of mHealth

Example prompts: What would prevent you from using mHealth systems?

Topic 4. Alerts and reminders

Example prompts: Is there any part of your/your patients' asthma management that is often forgotten?

Topic 5. User feedback and support

Example prompts: What type of support would you like? Examples include intuitive interfaces with information about asthma, FAQs, access to GP, specialist asthma nurse, speak to other users.

Topic 6. Privacy

Example prompts: How would you feel about personal medical data being stored on a mobile device/being data shared with your healthcare team and/or medical researchers?

Topic 7. Product design

Example prompts: What design aspects would you accept/find unacceptable? Would you consider carrying an additional device(s)?

groups. Because mHealth systems can provide multiple functions relevant to patients and healthcare professionals, we did not impose any restrictions on the number of responses that could be selected by participants and all responses were given equal weighting.

Individuals with asthma and healthcare professionals completed the questionnaires *via* an online survey platform (www.surveymonkey.com) over a 2-week period in December 2015. A hyperlink to the online survey was included on adverts placed online at Asthma UK and the European Commission websites and *via* the social media channels of Asthma UK and the European Federation of Allergy and Airways Diseases Patients' Associations (EFA). Healthcare professionals from the North West Severe Asthma Network and the North West respiratory postgraduate contact list were invited *via* e-mail to complete the online questionnaire. For purposes of characterising the sample, individuals with asthma completed the Asthma Control Questionnaire (ACQ); a score of ≥ 1.5 defined uncontrolled asthma and < 1.5 was classified as controlled or partly controlled asthma [21]. Chi-squared tests identified differences in response frequency between individuals with asthma and healthcare professionals and between individuals with controlled (including partly controlled) and uncontrolled asthma, using a statistical software package (SPSS, version 22.0). Significance was set at $p \leq 0.05$.

Data integration

Qualitative focus group data and quantitative results from the questionnaires were integrated under three core categories. Focus group quotes relating to questionnaire data were identified and used to illuminate and complement or contrast the quantitative results.

Ethics

A National Health Service (NHS) research ethics committee (15/EM/0360) and the ethics committee of Leiden University Medical Centre (P15.195) approved this study and participants gave informed consent.

Results**Participant characteristics**

The focus groups comprised 18 individuals with asthma (nine women) and five healthcare professionals (two medical doctors, two asthma nurses and a physiologist). The questionnaire was completed by 186 individuals with asthma (mean \pm SD age 40 ± 16 years, 135 women and 91 individuals with uncontrolled asthma) and 63 healthcare professionals (31 general practitioners, 13 hospital doctors, eight asthma nurses and 11 from other healthcare disciplines).

Core category 1: experiences and perceived uses of mHealth for asthma self-management

Individuals with asthma had considerably varied experience with mHealth, with some participants reporting no experience of using mHealth for their asthma and a few participants reporting considerable experience with multiple devices. Healthcare professionals' experience with mHealth systems was limited

to their use during research projects. The types of mHealth systems that participants had experience with are presented in table 2.

Focus group participants identified 12 potential uses of mHealth systems for asthma self-management (table 2). These proposed uses were developed into an item on each questionnaire, relating to 1) functions individuals with asthma would like from mHealth and 2) functions that healthcare professionals believed would be useful (table 3).

Patients most frequently requested an mHealth system to monitor asthma over time (72%) and to collect data to present to healthcare teams (70%) (table 3). This may relate to difficulties that patients have recalling symptoms and conveying these to their healthcare teams, as illustrated by quote 1.1 (table 4). Discussions with healthcare professionals revealed that they would find it useful if a patient presented them with data on an mHealth system, and one healthcare professional suggested that it might empower their patients (quote 1.2; table 4).

Healthcare professionals most frequently selected functions that alert patients to deteriorating asthma control (86%) and advise them when to seek medical attention (87%) (table 3). Focus group data highlighted that mHealth could prompt patients to seek medical attention sooner (quote 1.3; table 4). Support for these functions amongst individuals with asthma was also high (table 3) and comments in the focus group discussions were broadly aligned with those of the healthcare professionals (quotes 1.4 and 1.5; table 4).

A recurring theme in the focus group discussions was the potential to incorporate asthma action plans into mHealth. The preference of mHealth over the traditional pen and paper approach may be linked to the increased convenience/accessibility of mHealth (quote 1.6; table 4). In the questionnaires, 46% of patients and 79% of healthcare professionals ($p < 0.001$) answered that they would like or find it useful to have an asthma action plan incorporated into an mHealth system. The reason behind the greater support from healthcare professionals was not apparent from the focus group discussions.

Some of the proposed uses for mHealth systems that originated in the focus group discussions received less support from the questionnaires. In the focus groups, several patients voiced frustrations about attending routine asthma check-ups and proposed mHealth as a possible replacement (quote 1.7; table 4). However, in the surveys, only 25% of patients and 33% of healthcare professionals indicated that they would like or find it useful for an mHealth system to replace routine asthma check-ups.

Core category 2: useful measurements for managing asthma

The focus group discussions highlighted many measurements that participants believed could provide support for the self-management of asthma (table 2); these were developed into items on the questionnaires (table 5).

TABLE 2 Emergent themes identified in focus groups with people with asthma (n=18) and healthcare professionals (n=5) integrated under core categories

Emergent themes	
Core category 1	
Experience of mHealth	Experience with applications for nutrition analysis, inhaler (medication) monitoring, activity level monitoring, lung function (peak flow) monitoring, mental health, environmental monitoring (e.g. pollution and pollen), asthma diary.
Potential uses of mHealth	Replace check-ups, advise when to seek medical attention, monitor asthma over time, collect data to present to healthcare team, alerts to deterioration in asthma control, use as an asthma action plan, provide education materials, instructions on how to manage an asthma attack, a system to call for emergency help, a system to update medical records, a system to record side effects, a system to determine medication effectiveness.
Core category 2	
Useful measurements	Environment conditions (e.g. pollution, allergens (pollen), temperature), lung function (e.g. peak flow and measurements of airway inflammation), breathing (e.g. breathing rate and details of how often you cough), heart rate and activity levels, stress level, medication adherence, inhaler technique, diet, quality of sleep, self-reported symptoms.
Useful alerts	Medication running low, using their medication too much, they have not taken their inhaler, they are using their inhaler incorrectly, lung function is getting worse, pollution level in their area is high, pollen/allergen levels in their area are high, temperature/humidity in their area is high/low.
Core category 3	
Acceptability and barriers	Usability, product design, privacy, time, personalisability, human contact, data usefulness (e.g. reliability, interpretation and subjectivity), cost, mobile compatibility, medication compatibility.

TABLE 3 Questionnaire results: what individuals with asthma would like from an mHealth system and what healthcare professionals (HCPs) believe would be useful functions

Response options	Asthma [#]	HCPs [¶]	p-value ⁺	Asthma		p-value ^{##}
				Uncontrolled [§]	Controlled ^f	
A device/system that could replace routine (e.g. annual) asthma check-ups	25	33	0.21	12	40	<0.001
A device/system that offers advice regarding when additional medical attention should be sought	49	87	<0.001	56	44	0.12
A device/system to help patients monitor their asthma over time	72	81	0.14	77	66	0.12
A device/system to collect data that patients can show their doctor/healthcare professional, to demonstrate how their asthma has been	70	78	0.30	71	67	0.57
A device/system that detects and alerts patients and/or healthcare professionals to a deterioration in their asthma control before they would normally notice	69	86	0.01	75	64	0.18
A device/system for patients to use as their asthma action plan	46	79	<0.001	53	40	0.08
A device/system to offer educational materials about asthma	22	73	<0.001	25	17	0.21
A device/system that provides instructions on how to manage their asthma in an emergency	45	81	<0.001	47	44	0.68
A device/system that can be used to call for emergency help during an asthma attack	49	52	0.69	52	49	0.71
A device/system that can take measurements and update a patient's medical record	53	51	0.80	56	49	0.34
A device/system to record treatment side effects	44	37	0.29	46	44	0.79
A device/system that can tell if changes to patient's asthma medication have improved their asthma control	36	76	<0.001	45	28	0.02

Data are presented as percentages. Bold text indicates significant differences between groups. [#]: people with asthma (n=186); [¶]: HCPs, healthcare professionals (n=63); ⁺: patients with asthma *versus* HCPs; [§]: individuals with an asthma control questionnaire score ≥ 1.5 (n=91); ^f: individuals with an asthma control questionnaire score <1.5 (n=86); ^{##}: patients with controlled *versus* uncontrolled asthma.

Lung function measurements (71%) were commonly identified as being helpful to maintain asthma control (table 5). This was linked with patients' perception of a connection between asthma control and lung function and when to take appropriate action (quotes 2.1 and 2.2; table 4). Additional physiological parameters identified as being useful for asthma control included resting heart rate, breathing rate, stress levels, sleep quality and diet. These measurements were given varying, but usually modest, support from the questionnaire data (table 5).

Measurements regarding environmental conditions were believed to be helpful for asthma self-management by 71% of individuals with asthma and 68% of healthcare professionals (table 5). Focus group data suggest that environmental alerts may affect individuals' behaviours (quote 2.3; table 4).

Both individuals with asthma and healthcare professionals identified the negative impact of incorrect inhaler technique and provided support for their integration into mHealth (quotes 2.4 and 2.5; table 4). However, survey data highlighted a notable contrast in the results, with a significantly higher proportion of healthcare professionals compared with patients believing that measuring inhaler technique (87% *versus* 43%, $p<0.001$) and medication adherence (89% *versus* 48%, $p<0.001$) would be helpful for asthma control.

Core category 3: acceptability of and barriers to using mHealth systems for asthma self-management

Table 2 summarises perspectives on the acceptability of and barriers to the use of mHealth systems. The ability to personalise mHealth systems was a consistent subcategory across all focus groups. Discussions highlighted that different populations, e.g. children, the elderly and people with differing asthma severity, have different user requirements. Furthermore, it was proposed that mHealth systems may need to be personalised at an individual level (quote 3.1; table 4). One participant with asthma suggested that measuring numerous "irrelevant" parameters might discourage their compliance with mHealth (quote 3.2; table 4). Similarly, patients highlighted that if the burden of inputting data was too much then they would not be willing to comply with the device and emphasised that mHealth should be as automated as possible (quote 3.3; table 4).

The topic of data usefulness was common across all focus groups and included comments regarding the reliability of data, data subjectivity and the interpretation of data. Individuals with asthma highlighted concerns

TABLE 4 Selection of supporting quotes

Quote number	Selected quotes
Core category 1	
1.1	"It would be handy having an app so that you can monitor [asthma] yourself ... to show your consultant and respiratory nurse exactly how your asthma has been ... because often when they ask you can't remember." [Patient #7, London]
1.2	"It's a very powerful tool to be able to show [patients] the data and say this is what is happening ... rather than just saying you've got to keep taking your medication ... you are empowering them with their treatment." [HCP #3]
1.3	"A little bit of a prompt to say that at these levels maybe you should be seeking medical attention, this would be helpful because then they may attend the accident and emergency department a little bit sooner." [HCP #2]
1.4	"Things sometimes get worse and I don't necessarily notice them and, therefore, I let them get worse. It would be nice if I could monitor it and see trends in different things and address them." [Patient #4, London]
1.5	"I have had that moment, where you think at what point do I call an ambulance ... I would like to be able to hit a button and it says this is what you should be doing." [Patient #3, London]
1.6	"If you could have your asthma check-up and plug in your asthma action plan values into something ... that's a much better way of helping people stay in control than a piece of paper, that when they come back from the doctors they put down and don't touch again until the next asthma check." [Patient #3, Manchester]
1.7	"I really dislike going to my asthma check-up when I am pretty sure it is fairly well controlled anyway ... I go and they tell me what I already know ... it would be nice if a device could feed back to the nurse and they could let me know when I should get a check-up." [Patient #2, London]
Core category 2	
2.1	"My peak flow tends to go down and then I get worse ... even if I don't feel bad, my peak flow will be lower than it should be." [Patient #7, London]
2.2	"On the written asthma action plans, it says if my peak flow drops below 'X' then I should do this with my medication ... so it is useful for that." [Patient #3, London]
2.3	"If it's going to be a high pollen count, I will arrange to go out in the evening or very early in the morning and avoid that part of the day ... air quality is something that's a bit more difficult to avoid, but it's useful to know and may influence whether I go for a run today or whether I wait until tomorrow." [Patient #4, Manchester]
2.4	"The amount of times I rush through it [taking my inhaler] ... I don't feel like I am getting the benefit from it." [Patient #4, London]
2.5	"Patients are on step four or five treatment but can't take an inhaler correctly ... it is frightening that they are being referred to us for more invasive treatments." [HCP #1]
Core category 3	
3.1	"Not all of the asthmatic patients have the same symptoms ... I think you need to individualise the symptoms and what is measured to every patient separately." [HCP #5]
3.2	"I don't want to measure all those things ... if it is one or maybe two things I might, or if you could personalise it to what is relevant to you, but I'm not going to measure all of those things." [Patient #2, London]
3.3	"If it is automatically on your inhaler and it measures and gives you feedback, perfect ... if I have to get a separate device out to measure it, then I would probably use it less." [Patient #7, Leiden]
3.4	"I just don't like subjective questions. I don't remember how bad 'bad' was last time I selected bad." [Patient #2, London]
3.5	"If something is wearable and discreet, I would definitely go for something like that. If it is bulky and very visible, then maybe not." [Patient #4, London]
HCP: healthcare professional.	

with subjective measurements, such as self-reported symptoms (quote 3.4; table 4). The interpretation of the data, either by automated systems or by the individuals themselves, was highlighted as a source of possible error and an important consideration. Consequently, only 12% of patients responding to the questionnaire indicated that they would accept all recommendations to change their medication based on feedback from mHealth. This value increased to 30% if data supporting the recommendation was also presented to the patient and to 41% if the patient's doctor endorsed the mHealth system. Similarly, only 21% of healthcare professionals

TABLE 5 Questionnaire results: which of the following measurements do you think could help you/your patients achieve better asthma control?

Response options	Asthma [#]	HCPs [¶]	p-value ⁺	Asthma		p-value ^{##}
				Uncontrolled [§]	Controlled ^f	
Measurements of environment conditions (e.g. pollution, allergens, temperature and humidity)	70	68	0.81	75	65	0.16
Measurements of lung function (e.g. peak flow and measurements of airway inflammation)	71	75	0.58	71	70	0.82
Measurements of breathing (e.g. breathing rate and details of how often you cough)	64	60	0.60	68	60	0.29
Measurements of heart rate and activity levels	46	37	0.18	49	43	0.39
Measurements of stress levels	53	37	0.03	57	49	0.27
Measurements of medication adherence	48	89	<0.001	52	44	0.32
Measurements of inhaler technique	42	87	<0.001	43	43	0.98
Measurements of diet	32	32	0.94	36	24	0.09
Measurements of quality of sleep	54	44	0.20	58	48	0.16
Measurements of self-reported symptoms	34	57	<0.001	40	30	0.19

Data are presented as percentages. Bold text indicates significant differences between groups. [#]: people with asthma (n=186); [¶]: HCPs, healthcare professionals (n=63); ⁺: patients with asthma *versus* HCPs; [§]: individuals with an asthma control questionnaire score ≥ 1.5 (n=91); ^f: individuals with an asthma control questionnaire score <1.5 (n=86); ^{##}: patients with controlled *versus* uncontrolled asthma.

would be comfortable for their patients to change their medication based on mHealth feedback. This value increased to 46% if the healthcare professional could see patient data and approve the changes. Furthermore, 22% of healthcare professionals would like to see the patient in person before they recommended any changes to their medication.

Data security and data use was a common theme across focus groups. Patients expressed opposing views with regards to data security, with some patients unconcerned with how their data was managed and some insistent that data security is of utmost importance. Participants of the focus groups expressed unanimous support for their data to be used in an anonymous format for research purposes, whilst questionnaire results suggest just over half (58%) of patients were happy for anonymous data to be used for research purposes.

Physical properties relating to product design and compatibility were discussed as important considerations for mHealth. Questionnaire analysis revealed that 76% of individuals with asthma would be willing to carry or wear at least one additional device and 72% would be willing to keep an additional device at home. However, discussions in the focus groups indicate that this might depend on the product design (quote 3.5; table 4).

Discussion

The era of mHealth offers huge potential to enhance conventional healthcare. Asthma is an ideal candidate condition for mHealth developments, being a long-term condition that requires continuous attention from both healthcare professionals and patients. If mHealth systems are to be utilised in routine practice, they need to be embraced by both end-users. This is the first study to comprehensively explore patients' and healthcare professionals' perspectives on the use of mHealth for the self-management of asthma. There were significant differences in opinions with regards to expectations between healthcare professionals and patients; however, both end-users indicated substantial support for mHealth for asthma self-management.

All people with asthma should receive a personal asthma action plan as part of their asthma self-management strategy [7, 8]. However, only around a quarter of individuals with asthma receive such a plan [2, 11]. Our findings suggest that a large proportion of healthcare professionals believe that incorporating a personal asthma action plan into an mHealth system would be a useful function. This sentiment was commonly shared by individuals with asthma and vividly portrayed in the qualitative data, with one participant recalling the feeling of distress and indecision about how to treat their asthma and at what point to seek emergency attention. Neither patients nor healthcare professionals are enthusiastic about using written asthma action plans [9] and our data suggest the convenience of mHealth makes it an appealing alternative.

An accurate initial assessment and ongoing review of patients' asthma severity and control is crucial for the appropriate management of the disease [2]. Our data suggest that people find it difficult to express asthma severity and control to their healthcare teams, and indeed it is known that patients often underestimate their asthma severity [22, 23]. Such discrepancies between perception and objective asthma severity could have drastic consequences in the management of the disease. The UK National Review of Asthma Deaths suggests that poor recognition of asthma severity by patients and their healthcare teams,

and subsequent long-term under-treatment, are avoidable factors related to asthma deaths [2]. One proposed use of mHealth that was well supported by both end-users was a system that collects data over time, to assist patients in demonstrating their asthma control/severity to their healthcare teams. The selection of which parameters would be useful for this purpose merits careful consideration.

We identified a variety of physiological, environmental and behavioural measurements that individuals with asthma and healthcare professionals believe could support asthma self-management. Individuals with asthma most commonly responded that measurements of lung function would be useful for maintaining asthma control. The success of traditional asthma self-management programmes, relying on regular peak flow measurements [5], would support their belief. Other physiological parameters that were identified as being useful included measurements of heart rate, respiratory rate and sleep quality. Given that heart rate variability may be associated with asthma control [24], that respiratory rate variability during sleep may differ between individuals with and without asthma [25], and that nocturnal waking is a common complaint of individuals with asthma, the potential for these measurements to provide support for asthma self-management warrants further investigation.

Each year 5.5 million deaths can be attributed to poor air quality [26], whilst air pollution exposure is associated with an increased frequency of asthma attacks in children and adults [27]. A large proportion of individuals with asthma and healthcare professionals responding to our survey believe measurements of environmental conditions could help achieve better asthma control and should be incorporated into mHealth.

Asthma is no longer seen as a single disease, but a syndrome with heterogeneous presentation and numerous phenotypes and endotypes [28]. Participants identified that the complex and heterogeneous nature of asthma means that individuals will have differing requirements from mHealth. Whilst it is unrealistic to expect healthcare professionals to be trained in the use of multiple different mHealth systems, it was proposed by our participants that patients and their healthcare teams should be able to customise a panel of relevant functions and parameters for each patient. This poses a complex and challenging problem for mHealth developers, who should work in close partnership with a range of patients, with different levels of asthma control, and with healthcare professionals to ensure all end-user requirements are met.

Sharp contrasts were noted in the support for some mHealth functions between patients and healthcare professionals. Intriguingly, the functions that received less support from patients appear to relate to aspects that patients may be inherently aware of and therefore see no need for mHealth feedback, *e.g.* measurements of medication adherence, inhaler technique and self-reported symptoms. On the contrary, the well-supported functions relate to aspects whereby the patient would be somewhat blind to the information without such feedback, *e.g.* environmental and lung function measurements. It would seem a reasonable interpretation that patients advocate functions that relate to acquiring information that would otherwise be unknown to them, not simply monitoring parameters that they could already be aware of. In contrast, functions that include “big brother” monitoring of patients, such as medication adherence and inhaler technique, were well supported by healthcare professionals. This is reasonable given that healthcare professionals need to know that medication has been taken as prescribed in order to assess treatment efficacy, and adherence to treatment is known to be variable [29].

This research benefits from a mixed methods design, permitting themes identified in the focus groups to be quantified in the survey and integrated in the analysis to acquire a comprehensive understanding of the perspectives of patients and healthcare professionals on mHealth for asthma self-management. However, several methodological limitations deserve consideration. Participants’ responses to the questionnaires were given equal weighting in the analysis. This method fails to take into account the strength of their opinions. This study may therefore have benefited from the ability for participants to rank their responses in order of preference. This study may have also benefited from another round of focus group discussions, to probe further into the results from the questionnaire. The majority of the participants in the survey likely came from those who visit Asthma UK’s website or follow Asthma UK social media channels and, therefore, are likely to be more active and well-educated in the management of their asthma. The possibility of selection bias should therefore be considered during any interpretation of the results.

In conclusion, asthma is an ideal candidate for mHealth developments and recent times have seen a meteoric, but rather haphazard and often ill-informed [30], rise in mHealth systems for asthma self-management. A user-centred design of mHealth is integral for technology to meet end-users’ expectations and may improve adherence and health outcomes. This research provides overwhelming support for mHealth to assist asthma self-management, by both individuals with asthma and healthcare professionals, but highlights fundamental differences in preferred functions between the different end-users. It also identified numerous factors that would need consideration during the development of new mHealth devices. Developers of new mHealth systems should consider these opinions during the development of new user-centred mHealth systems to aid the self-management of asthma.

Acknowledgements

We would like to thank Joke De Vocht for her support in organising the patient advisory group and reviewing the focus group topic guides. We would also like to thank the members of the patient advisory group for their valuable insight and guidance in developing the topic guides.

Authors' contributions: A.J. Simpson (guarantor), P.J. Honkoop, E. Kennington, C. Coleman, K.F. Chung, J.K. Sont, O. Usmani and S.J. Fowler were responsible for the conception and design of the research; A.J. Simpson, E. Kennington, J.B. Snoeck-Stroband, I. Smith and J. East planned and conducted the focus groups; A.J. Simpson, J. East, E. Kennington and S.J. Fowler produced the questionnaire; A.J. Simpson and S.J. Fowler analysed results; A.J. Simpson, S.J. Fowler and A. Caress interpreted the results; A.J. Simpson drafted the manuscript; A.J. Simpson, P.J. Honkoop, E. Kennington, J.B. Snoeck-Stroband, I. Smith, J. East, C. Coleman, A. Caress, K.F. Chung, J.K. Sont, O. Usmani and S.J. Fowler edited and revised the manuscript; A.J. Simpson, P.J. Honkoop, E. Kennington, J.B. Snoeck-Stroband, I. Smith, J. East, C. Coleman, A. Caress, K.F. Chung, J.K. Sont, O. Usmani and S.J. Fowler approved the final version of the manuscript. Transparency declaration: A.J. Simpson had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. A.J. Simpson affirms that the manuscript is an honest, accurate and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned have been explained.

References

- Masoli M, Fabian D, Holt S, *et al.* The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 2004; 59: 469–478.
- Levy M, Andrews R, Buckingham R, *et al.* Why asthma still kills: the National Review of Asthma Deaths (NRAD) confidential enquiry report. Royal College of Physicians 2014. www.rcplondon.ac.uk/projects/outputs/why-asthma-still-kills Date last accessed: October, 2016.
- Powell H, Gibson PG. Options for self-management education for adults with asthma. *Cochrane Database Syst Rev* 2003; 1: CD004107.
- Gibson PG, Powell H. Written action plans for asthma: an evidence-based review of the key components. *Thorax* 2004; 59: 94–99.
- Gibson PG, Coughlan J, Wilson AJ, *et al.* Self-management education and regular practitioner review for adults with asthma. *Cochrane Database Syst Rev* 2000; 2: CD001117.
- Adams RJ, Smith BJ, Ruffin RE. Factors associated with hospital admissions and repeat emergency department visits for adults with asthma. *Thorax* 2000; 55: 566–573.
- British Thoracic Society, Scottish Intercollegiate Guidelines Network (SIGN). British guideline on the management of asthma; a national clinical guideline 2014. www.brit-thoracic.org.uk/document-library/clinical-information/asthma/btssign-asthma-guideline-2014/ Date last accessed: October, 2016.
- Global Initiative for Asthma (GINA). Global Strategy for Asthma Management and Prevention (2016 update). http://ginasthma.org/wp-content/uploads/2016/04/GINA-2016-main-report_tracked.pdf Date last accessed: October, 2016.
- Jones A, Pill R, Adams S. Qualitative study of views of health professionals and patients on guided self management plans for asthma. *BMJ* 2000; 321: 1507–1510.
- Honkoop PJ, Taylor DR, Smith AD, *et al.* Early detection of asthma exacerbations by using action points in self-management plans. *Eur Respir J* 2013; 41: 53–59.
- Centers for Disease Control and Prevention (CDC). Asthma facts; CDC's National Asthma Control Program grantees, 2013. www.cdc.gov/asthma/pdfs/asthma_facts_program_grantees.pdf Date last accessed: October, 2016.
- Kaya Z, Erkan F, Ozkan M, *et al.* Self-management plans for asthma control and predictors of patient compliance. *J Asthma* 2009; 46: 270–275.
- van Gaalen JL, Beerthuis T, van der Meer V, *et al.* Long-term outcomes of internet-based self-management support in adults with asthma: randomized controlled trial. *J Med Internet Res* 2013; 15: e188.
- Wu AC, Carpenter JF, Himes BE. Mobile health applications for asthma. *J Allergy Clin Immunol Pract* 2015; 3: 446–448.
- Kikidis D, Konstantinos V, Tzovaras D, *et al.* The digital asthma patient: the history and future of inhaler based health monitoring devices. *J Aerosol Med Pulm Drug Deliv* 2016; 29: 219–232.
- Marciano Belisario JS, Huckvale K, Greenfield G, *et al.* Smartphone and tablet self management apps for asthma. *Cochrane Database Syst Rev* 2013; 11: CD010013.
- McCurdie T, Taneva S, Casselman M, *et al.* mHealth consumer apps: the case for user-centered design. *Biomed Instrum Technol* 2012; Suppl., 49–56.
- Creswell JW, Clark VLP. Designing and Conducting Mixed Methods Research. London, SAGE publications Ltd, 2007.
- Greenbaum TL. The Handbook for Focus Group Research. 2nd Edn. London, SAGE Publications Ltd, 1997.
- Ritchie J, Spencer L. Qualitative data analysis for applied policy research. In: Bryman A, Burgess RG, eds. *Analyzing Qualitative Data*. Oxford, Routledge, 2002; pp. 173–194.
- Juniper EF, O'Byrne PM, Guyatt GH, *et al.* Development and validation of a questionnaire to measure asthma control. *Eur Respir J* 1999; 14: 902–907.
- Lurie A, Marsala C, Hartley S, *et al.* Patients' perception of asthma severity. *Respir Med* 2007; 101: 2145–2152.
- Nguyen BP, Wilson SR, German DF. Patients' perceptions compared with objective ratings of asthma severity. *Ann Allergy Asthma Immunol* 1996; 77: 209–215.
- Lutfi MF. Autonomic modulations in patients with bronchial asthma based on short-term heart rate variability. *Lung India* 2012; 29: 254–258.
- Campana LM, Owens RL, Butler JP, *et al.* Variability of respiratory mechanics during sleep in overweight and obese subjects with and without asthma. *Respir Physiol Neurobiol* 2013; 186: 290–295.
- GBD 2013 Risk Factors Collaborators, Forouzanfar MH, Alexander L, Anderson HR, *et al.* Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015; 386: 2287–2323.

- 27 Künzli N, Kaiser R, Medina S, *et al.* Public-health impact of outdoor and traffic-related air pollution: a European assessment. *Lancet* 2000 356: 795–801.
- 28 Skloot GS. Asthma phenotypes and endotypes: a personalized approach to treatment. *Curr Opin Pulm Med* 2016; 22: 3–9.
- 29 Lindsay JT, Heaney LG. Nonadherence in difficult asthma – facts, myths, and a time to act. *Patient Prefer Adherence* 2013; 7: 329–336.
- 30 Huckvale K, Car M, Morrison C, *et al.* Apps for asthma self-management: a systematic assessment of content and tools. *BMC Med* 2012; 10: 144.